

Varicella Outbreak Involving Two Dose Vaccine Recipients — Arkansas, 2006

Adriana S. Lopez, MHS

Division of Viral Diseases

**National Center for Immunization and Respiratory
Diseases**

Centers For Disease Control and Prevention

June 28, 2007



SAFER • HEALTHIER • PEOPLE™



New ACIP Recommendations for Varicella Immunization

- June 2005
 - ◆ 2nd dose for children to control outbreaks, resources permitting
- June 2006
 - ◆ Routine 2-dose vaccination for all children
 - ★ 1st dose: 12 – 15 months
 - ★ 2nd dose: 4 – 6 years
 - ★ Catch-up: >6 years

Arkansas Outbreak: September 2006

- 31 cases (as of 10/2)
 - ◆ 8 were 2-dose vaccine recipients
 - ◆ All reported mild or atypical disease
 - ◆ Elementary school complex (School A (pre-K), School B (K-3), School C (4-6))
 - ★ Prior outbreak in same complex January 2006
 - ★ Vaccination Clinic (per ACIP): ~ 400 students
 - Many received 2nd dose
- CDC invited to assist with current outbreak investigation
 - ◆ Objectives:
 - ★ Confirm varicella in 2-dose vaccine recipient
 - ★ Characterize vaccine effectiveness among 1- and 2-dose recipients

Outbreak Case Definition

- CSTE Case definition (acute maculopapulovesicular rash without other apparent cause) occurring between Sept 1 and Dec 18
 - ◆ In vaccinated persons disease may be:
 - ★ Mild
 - ★ Fewer than 50 skin lesions
 - ★ Shorter duration of illness
 - ★ Atypical in appearance (maculopapular with few or no vesicles)
- Operationally used 3 lesions as lower cut-off
- Lab confirmation
 - ◆ PCR (lesions, environmental samples)
 - ◆ IgM (blood, saliva)

Survey and Case Investigation

- Cases in students identified from:
 - ◆ Health Unit
 - ◆ School nurse
 - ◆ School wide survey:
 - ★ Vaccination status & varicella history
 - ★ Medical conditions
 - ★ “Rashes, insect bites, bumps, spots, or blisters since start of school year”
- Case investigation
 - ◆ Clinical information
 - ◆ Medical conditions
 - ◆ Medications

Vaccination Records


- Arkansas Immunization Registry
 - ◆ Vaccination status
 - ◆ Limited disease history (on unvaccinated cases)
- Alternate sources
 - ◆ Local paper records
 - ◆ Parental survey
- Vaccinations ≤ 42 days prior to rash onset
 - ◆ 1st dose = Unvaccinated
 - ◆ 2nd dose = Indeterminate

School-wide survey

- Overall response rate: 79%
 - ◆ School A (pre-K): 81%
 - ◆ School B (K-3): 84%
 - ◆ School C (4-6): 72%†
- Responders vs non responders
 - ◆ No difference in:
 - ★ Gender
 - ★ Vaccination status
 - ◆ Difference in:
 - ★ Race/ethnicity – African-American children less likely to be responders (65% vs. 86% for white/Hispanic, $p < 0.001$)

† $p < 0.001$

Reported varicella cases by rash onset date and vaccination status -- School Complex, Arkansas, 9/1-12/18/2006 (n=83*)



*2 cases with unknown rash onset dates excluded

Disease Severity

| | No Prior Disease | | Prior disease (n=15) | P-value |
|--------------------|------------------|------------------|-------------------------|---------|
| | 1 dose (n=47) | 2 dose (n=23) | | |
| <hr/> | | | | |
| Number of lesions* | | | | |
| <50 | 81% (38) | 91% (21) | 67% (10) | 0.20 |
| 50-249 | 13% (6) | 0% (0) | 7% (1) | |
| 250-499 | 0% (0) | 0% (0) | 0% (0) | |
| ≥500 | 0% (0) | 0% (0) | 0% (0) | |
| <hr/> | | | | |

*Number of lesions was unknown for 3 cases with 1-dose, 2 cases with 2-doses, and 4 cases with previous disease.


Disease Characteristics

| | No Prior Disease | | Prior disease (n=15) | P-value |
|-------------------------------------|------------------|------------------|-------------------------|---------|
| | 1 dose (n=47) | 2 dose (n=23) | | |
| Rash description | | | | |
| Maculopapular | 83% (39) | 83% (19) | 60% (9) | 0.10 |
| Vesicular | 45% (21) | 43% (10) | 33% (5) | 0.49 |
| Itchy | 87% (41) | 87% (20) | 100% (15) | 0.23 |
| Median rash duration, days (range)* | 6 (1-13) | 4 (1-16) | 4 (1-7) | 0.37 |
| Fever | 30% (14) | 17% (4) | 20% (3) | 0.31 |

*Based on available data for 77 cases

Case Example Highlighting Challenges to Case Ascertainment

- Clinical characteristics of a 2 dose vaccine recipient
 - ◆ Only 4 papules at any one time
 - ◆ 1 to 2 vesicles
 - ◆ Itchy
 - ◆ Lasted 3 – 4 days
 - ◆ Specimens obtained on day of rash onset
 - ★ Lesion: PCR positive
 - ★ Saliva: Negative



Vaccination and Disease History of Cases (N=85)

| Vaccination Status | Cases (%) | Attack rate |
|---|-------------------|-------------|
| 2 doses <i>- w/prior disease</i> <i>- w/unknown disease hx</i> | 25 (29) 2 1 | 10.4 |
| 1 dose <i>- w/prior disease</i> | 54 (64) 7 | 14.6 |
| 0 doses <i>- w/prior disease</i> | 6 (7) 6 | -- |

Vaccination Coverage

| VACCINATION STATUS | SCHOOL | | | TOTAL N = 758* |
|--------------------|-----------------|------------------|------------------|-------------------|
| | PRE-K N = 98 | K – 3 N = 395 | 4 – 6 N = 265 | |
| Unvaccinated | 1% | 2% | 4% | 3% |
| 1 Dose | 84% | 54% | 48% | 56% |
| 2 Doses** | 15% | 44% | 48% | 41% |
| Total Vaccinated | 98% | 97% | 96% | 97% |

*Excluded:

106 students with history of disease

9 students with no vaccination data

** Includes 22 cases with 2nd dose within 42 days

Laboratory Results

- 27 cases tested: 6 positive
 - ◆ 5 PCR (lesion)
 - ★ 1 two dose recipient
 - ★ 3 one dose recipients
 - ★ 1 unvaccinated with history of disease
 - ◆ 1 IgM serum spot (one dose recipient)
- Environmental samples from school and case-patient bedding
 - ◆ Pajamas and pillowcase PCR positive for wild-type virus

Vaccine Effectiveness Against Clinical Varicella Using Historic Attack Rates

- If we compare attack rates with historic rates for any varicella disease in unvaccinated:
 - ◆ AR_{unvacc} of 80%
 - ★ 2 doses to unvaccinated:
$$VE_2 = 1 - (10.4\% / 80\%)$$
$$= \mathbf{87\% (95\% CI: 80.5 to 91.4\%)}$$
 - ★ 1 dose to unvaccinated:
$$VE_1 = 1 - (14.6\% / 80\%)$$
$$= \mathbf{81.8\% (95\% CI: 75.8 to 86.2\%)}$$

Limitations

- May not have had sufficient power to detect differences
- Misclassification of case status
 - ◆ Broad clinical definition → False positives
 - ◆ Mild disease → False negatives
 - ◆ Lesion-based laboratory diagnostics only helpful while transient rash still present
- Misclassification of disease history
 - ◆ Based on parental report
- Response rate
 - ◆ Lower among School C students

Summary

- Varicella confirmed as cause of outbreak on both epidemiologic and laboratory criteria
 - ◆ Lab confirmation included two-dose recipient
- 85 cases identified
 - ◆ 25 (29%) two-dose recipients
 - ◆ Similar clinical picture in 1 vs. 2 dose recipients
- 97% school vaccination coverage
 - ◆ 41% two doses*
- 2-dose VE point estimate 5% points higher than 1-dose VE but overlapping confidence intervals

*Includes persons with 2nd dose < 42 days

Conclusions

- One of the largest varicella outbreaks investigated in recent years
- First U.S. outbreak reported with significant number of second dose vaccine recipients
- Case ascertainment challenging and may preclude evaluation of 2 vs. 1 risk reduction assessments in outbreak settings
- Moderate 2-dose coverage was insufficient to prevent this outbreak

Next steps

- Additional vaccine effectiveness studies needed
 - ◆ Range of VE for 1-dose: 44-100% from >30 studies
- Need to monitor the number and size of outbreaks as a key outcome to assess effects of the routine 2 dose policy

Acknowledgements

- Ashley County Health Unit- Hamburg
 - ◆ Connie Scott
 - ◆ Nan Erwin
- Southeast Public Health Region Office
 - ◆ Donetta McGriff
 - ◆ Carolyn (Sue) Hutton
- AR State Health Department
 - ◆ Linda Gladden
 - ◆ Karen Fowler
 - ◆ Haytham Safi
 - ◆ Sandra Snow
 - ◆ Helen Dang
 - ◆ AR State Lab
- Schools
- CDC Atlanta
 - ◆ Philip Gould
 - ◆ Jessica Leung
 - ◆ Meredith Reynolds
 - ◆ Sandra Chaves
 - ◆ Rafael Harpaz
 - ◆ Jane Seward
 - ◆ Tureka Watson
 - ◆ John Zhang
 - ◆ Scott Schmid
 - ◆ National VZV Lab

Thank you

The findings and conclusions in this presentation have not been formally disseminated by the Centers for Disease Control and Prevention and should not be construed to represent any agency determination or policy



SAFER • HEALTHIER • PEOPLE™

